

**FIG.\_1****FIG.\_2**

2 / 3



STEP 1: SET UP 3 PCR REACTIONS:


PRODUCTS:

TUBE 1: 

TUBE 2: 


TUBE 3: 

STEP 2: SET UP PCR REACTION WITH PRODUCTS OF TUBE 1 + PRODUCTS TUBE 2 + F1 + R4.





STEP 3: REPEAT STEP 2 USING PRODUCT FROM STEP 2 + PRODUCT FROM STEP 1, TUBE 3 + PRIMERS F1 + R3.


**FIG.\_3**



STEP 1: SET UP 3 PCR REACTIONS:


TUBE 1:   
RE1

TUBE 2:   
RE2


TUBE 3:   
RE2

STEP 2: DIGEST PRODUCTS FROM STEP 1 WITH SUITABLE RESTRICTION ENDONUCLEASES.


STEP 3: LIGATE DIGESTED PRODUCT FROM STEP 2, TUBE 2 WITH DIGESTED PRODUCT FROM STEP 2, TUBE 1.




STEP 4: AMPLIFY VIA PCR LIGATED PRODUCTS OF STEP 3 WITH F1 + R4.



STEP 5: DIGEST AMPLIFIED PRODUCT OF STEP 4 WITH RESTRICTION ENDONUCLEASE #2.




STEP 6: LIGATE PRODUCT FROM STEP 5 WITH PRODUCT FROM STEP 2, TUBE 1.



STEP 7: AMPLIFY PRODUCT FROM STEP 6 WITH F1 + R3.

**FIG.\_4**



**FIG.\_5**